

# Genetic Aspects of Hepatocellular Carcinogenesis

MEHMET OZTURK, Ph.D

**ABSTRACT:** *Hepatocellular carcinoma (HCC) is linked etiologically to viruses (hepatitis B virus [HBV] and hepatitis C virus [HCV]), chemical carcinogens (i.e., aflatoxins), and other environmental and host factors causing chronic liver injury. Some hepatoblastomas may be linked to inherited gene mutations, but adult hereditary HCC appears to be rare. HCCs display gross genomic alterations, including DNA rearrangements associated with HBV DNA integration, loss of heterozygosity, and, less importantly, chromosomal amplifications and loss of imprinting. Many genes with somatic mutations have now been identified in these tumors. Most frequently involved genes are tumor suppressor genes such as p53, M6P/IGF2R,  $\beta$ -catenin, p16INK4A, and retinoblastoma genes. Most identified mutations are somatic, but germline mutations of p16INK4A, APC, and BRCA2 have also been reported. Oncogenic activation of several cellular genes such as cyclin D and cyclin A have been described in HCC, but the possible implication of candidate viral oncogenes (i.e., X protein of HBV) is still debated. A comprehensive analysis of all the genetic changes described for HCC demonstrates that at least four different growth regulatory pathways are altered in these tumors. However, each pathway appears to be implicated in a limited fraction of these tumors, suggesting that HCCs are genetically heterogeneous neoplasms. This genetic heterogeneity correlates with the heterogeneity of etiologic factors implicated in HCC.*

**KEY WORDS:** hepatocellular carcinoma, primary liver cancer, p53, p16INK4A, cyclin D,  $\beta$ -catenin, M6P/IGF2R

Hepatocellular carcinoma (HCC) cells often display chromosomal changes such as polyploidy, loss of heterozygosity (LOH), allelic imbalance (AI), amplifications, and translocations.<sup>1</sup> It has also been known for a long time that hepatitis B virus (HBV) DNA causes chromosomal rearrangements by integration into the host genome.<sup>2</sup> It is expected that the chromosomal regions that undergo tumor-specific changes harbor critical genes involved directly (oncogenes and tumor suppressor genes) or indirectly (DNA repair genes) in carcinogenesis. To date, a dozen genes, including p53, mannose-6-

phosphate/insulin-like factor 2 receptor (M6P/IGF2R),  $\beta$ -catenin, retinoblastoma (Rb1), p16INK4A, adenomatosis polyposis coli (APC), breast cancer gene 2 (BRCA2), cyclin A, cyclin D, and insulin-like growth factor 2 (IGF2) have been shown to be altered in HCC and/or hepatoblastoma (Table 1). This list will probably grow over the next years to include many more genes. There are at least two reasons to explain the high number of altered genes in HCC. First, solid tumors of the adult may need the accumulation of many genetic alterations before they become clinically detectable. Indeed,

## Objectives

Upon completion of this article, the reader should be able to: 1) list the factors that are etiologically linked to hepatocellular carcinoma; 2) state the most frequently involved genes; and 3) recognize the four different growth regulatory pathways that are altered in these tumors.

## Accreditation

The Indiana University School of Medicine is accredited by the Accreditation Council for Continuing Medical Education to sponsor continuing medical education for physicians.

## Credit

The Indiana University School of Medicine designates this educational activity for a maximum of 1.0 hours credit toward the AMA Physicians Recognition Award in category one.

## Disclosure

Statements have been obtained regarding the author's relationships with financial supporters of this activity. There is no apparent conflict of interest related to the context of participation of the author of this article.

From the Department of Molecular Biology and Genetics, Bilkent University, 06533 Ankara, Turkey.

Reprint requests: Dr. Mehmet Ozturk, Dept. of Molecular Biology and Genetics, Bilkent University, 06533 Ankara, Turkey.

Copyright © 1999 by Thieme Medical Publishers, Inc., 333 Seventh Avenue, New York, NY 10001, USA. Tel.:

+1(212) 760-0888 x132.0272-8087/1999/1098-8971 (1999) 19:03:0235-0242:SLD00010X

**TABLE 1. Genetic Alterations in Hepatocellular Carcinoma and Hepatoblastoma**

<i>Gene</i>	<i>Mutation (%)</i>	<i>Other Alterations</i>	<i>References</i>
<i>p53</i>	28	HBx interaction	see Table 2
<i>M6P/IGF2R</i>	18–33	HD	21, 36, 44
<i>TGFRB2</i>	0	—	23
<i>RB1</i>	15	LOH, HD	11,‡ 45–50
<i>p15INK4B</i>	0	HD	50, 51
<i>p16INK4A*</i>	0–55	HD, Methylation	50–54
<i>p21</i>	5	—	55
Cyclin D <sup>†</sup>	11–13	—	56, 57
Cyclin A <sup>†</sup>	19	HBV integration	58, 59
β-catenin	19–26	—	30, 31
<i>APC*</i>	62‡	LOH	11, 32–35, 37
E-cadherin	NR	LOH, Methylation	60, 61
<i>BRCA2*</i>	5	LOH	62
<i>Smad2</i>	0–2	—	24, 25
<i>Smad4</i>	0–6	—	24, 25
<i>hMLH1</i>	NR	LOH	63
<i>hMSH2</i>	NR	LOH	63
<i>IGF2</i>	NR	LOI	38–43
<i>K-ras</i>	0–17	—	90–95
<i>N-ras</i>	0–16	Amplification	90, 91, 93, 95, 96
<i>H-ras</i>	0–10	Methylation	73, 91, 93, 95, 97, 98
<i>c-myc</i> <sup>†</sup>	0–50	—	99, 100
<i>N-myc</i> <sup>†</sup>	0	—	99, 101

\*Somatic and germline mutations.

<sup>†</sup>Amplification.

‡In hepatoblastoma only.

HD, homozygous deletion.

LOH, loss of heterozygosity.

LOI, loss of imprinting.

the well-known “latent period” between the first exposure to an etiologic agent (i.e., infection with HBV) and the development of HCC is in favor of such a hypothesis.<sup>3</sup> Second, the multiplicity of genetic alterations in HCC may indicate that different etiologic factors affect different sets of target genes in hepatocytes. This etiologically defined genetic heterogeneity of HCC results in a phenotypic heterogeneity of these tumors. In other words, distinct but related growth regulatory pathways are altered during hepatocarcinogenesis. As discussed later, at least four different pathways are altered in human HCCs.

### ***p53* GENE**

Many reports now indicate that the *p53* gene, which is located on chromosome 17p, is mutated in about 30% of HCCs worldwide (for a recent review, see ref. 4). All reported *p53* mutations in HCC are somatic. Therefore, germline mutations of *p53* appear not to predispose to HCC. Both the frequency and the type of *p53* mutations are different depending on geographic location and suspected etiology of these tumors (Table 2). An HCC-specific codon 249 mutation (AGG → AGT) leading to an arginine to serine substitution (R249S), suspected to be induced by aflatoxins, was found in most HCCs from geographic areas with high incidence of HCC and a

high risk of exposure to aflatoxins.<sup>5–7</sup> This mutation was found in 50% of HCCs from Mozambique,<sup>8,9</sup> 50 to 75% of HCCs from Qidong province of China,<sup>6,10,11</sup> and 67% of HCCs from Senegal.<sup>7</sup> A worldwide study by Ozturk et al.<sup>8</sup> suggested a close correlation between the presence of codon 249 mutations in HCC and high risk of aflatoxin intake. This early study has now been largely confirmed by others. As shown in Table 1, the codon 249 mutation is present in 36% of tumors from Africa and 32% of tumors from China, respectively. These two regions of the world are known for high incidence of HCC, where both HBV and aflatoxins are recognized as major etiologic factors. In contrast, the codon 249 mutation is seen in less than 4% of HCCs from Japan, Europe, and North America, where HBV and hepatitis C virus (HCV), but not aflatoxins, are the main etiologic factors. The overall frequency of codon 249 mutations in the world is 11%. Mutations affecting other codons of the *p53* gene are detected in HCC, and their worldwide frequency is 18%. The frequency of all *p53* mutations in HCC varies between 15% in Europe and 42% in China, with a worldwide frequency of 27% (see Table 2 for a detailed analysis of *p53* mutations). Thus, *p53* gene is mutated in about a third of HCCs, but only a third of these mutations can be etiologically linked to a high risk of aflatoxin exposure. Therefore, *p53* mutations can occur in HCC independent of aflatoxin risk, and in HBV or HCV infection. However, Unsal et al.<sup>9</sup> reported an

**TABLE 2. Frequency of *p53* Mutations in Hepatocellular Carcinoma**

Region	<i>p53</i> -249 <sub>ser</sub>	<i>p53</i> -other	<i>p53</i> -total
North America	4/89* (4%)	6/34 (18%)	6/34 (18%)
South America†	3/16 (19%)	ND —	ND —
Europe	0/131 (0%)	13/109 (12%)	23/151 (15%)
Africa‡	19/53 (36%)	2/26 (8%)	11/26 (42%)
China§	62/191 (32%)	16/132 (12%)	50/132 (38%)
Japan	7/292 (2%)	53/232 (23%)	94/353 (27%)
Other	20/265 (8%)	57/265 (22%)	79/273 (29%)
Total	115/1037 (11%)	147/798 (18%)	263/969 (27%)

Data were compiled from references 6, 7, 9–11, 45, 64–89, 98 and 102–104. The numbers do not add up for two reasons: in some studies only the *p53*-249 mutation was reported; in some others only the information of the total number of mutations was reported.

\*U.S.A. including Alaska.

†Mexico only.

‡South Africa, Mozambique, and Senegal only.

§Mainland China and Hong Kong.

||Australia, Singapore, South Korea, Taiwan, Thailand.

apparent association between the presence of X gene coding sequences of HBV (HBx) and wild-type *p53* in HCC. Based on this observation, a possible interference of HBV with wild-type *p53* function was suggested.<sup>9</sup> Indeed, recent studies showed that HBx protein encoded by the X region of HBV interacts with wild-type *p53* protein both physically and functionally.<sup>12–15</sup> These observations suggest that the suspected oncogenic activity of HBx protein is linked to functional inactivation of wild-type *p53* protein, as observed with other viral proteins with transforming activity. However, the interaction of HBx with *p53* was shown only experimentally. It is presently unclear whether HBx-*p53* interactions really occur in HBV-infected hepatocytes and/or in HCC cells with integrated HBV DNA sequences.

Frequent involvement of *p53* mutations in HCC is not surprising for several reasons. First, the *p53* gene is the only known gene to be mutated at a very high frequency in tumors of different origin.<sup>16</sup> Second, this protein is involved in different cellular processes (cell cycle arrest, apoptosis, differentiation, angiogenesis, etc.), all critically involved in the development of malignancy.<sup>17</sup> Under physiologic conditions, *p53* protein is complexed with MDM2 protein that promotes a rapid degradation of *p53*. MDM2-*p53* complexes are inhibited either by p19ARF (induced by both cellular and viral oncogenes) or by N-terminal phosphorylation of *p53* by DNA-dependent protein kinase. This leads to an accumulation and functional activation of *p53* in cells, leading to either cell cycle arrest by p21 or apoptosis by

*bax* induction. Thus, *p53* protein appears to be involved in a growth control response to abnormal oncogene expression and DNA damage.<sup>17</sup> In patients with chronic liver disease, the risks of oncogene activation and DNA damage are elevated. As stated earlier, HBx may have an oncogenic activity and aflatoxins are potent DNA damaging agents.<sup>1,2</sup>

### ***p16INK4A*, CYCLIN D, AND RETINOBLASTOMA GENES**

These three genes encode for proteins involved in the regulation of the G1 phase of the cell cycle. Cyclin D forms active complexes with CDK4 protein, whereas *p16* protein is an inhibitor of CDK4 activity.<sup>18</sup> The retinoblastoma protein (pRb) is the main known substrate of CDK4. In nonproliferating cells, pRb protein forms complexes with E2F transcription factors. When complexed to pRb, E2Fs are transcriptionally inactive. Upon phosphorylation by CDK4, pRb is released from its complexes and “free E2Fs” promote the initiation of DNA synthesis.<sup>19</sup> These observations predict that the loss of pRb protein or its aberrant phosphorylation will lead to a loss of growth control at the G1 phase of the cell cycle. Increased phosphorylation of pRb may result from an aberrant activation of CDK4 by either an excess of cyclin D and/or a deficit in *p16* protein. Recent studies demonstrated that all three genes, namely *RB1*,

*p16INK4A*, and cyclin D, undergo structural alterations in HCC. The retinoblastoma gene (*RB1*) is one of the tumor suppressor genes studied in HCC just after the implication of *p53* in these tumors. LOH at the *RB1* gene locus is quite frequent in HCC. In addition, *RB1* mutations were observed in 15% of these tumors (Table 1).


The *p16INK4A* gene, which is located at chromosome 9p, codes for two alternatively spliced transcripts.<sup>17,18</sup> One of the transcripts is for p16 protein, an inhibitor of cyclin-dependent kinases 4 and 6.<sup>18</sup> *p16INK4A* status in HCC has been studied independently by several laboratories. Both germline and somatic mutations of *p16INKA* were found in HCC patients. It was also reported that about 50% of HCC display *de novo* methylation of *p16INK4A*, as observed in other cancers (see Table 1 and references therein). It is known that *de novo* methylation is a mechanism involved in gene silencing.<sup>20</sup> Therefore, one can assume that HCC cells with methylated *p16INK4A* are unable to express the gene, leading to the loss of a cyclin-dependent kinase inhibitor protein.

As shown in Table 1, cyclin D and cyclin A genes were shown to be amplified in 10–20% of HCCs. It is noteworthy that *RB1*, *p16INKA*, and cyclin genes are mutated *individually* in 10 to 20% of HCCs. Although

this frequency is not high, their involvement in the same growth regulatory pathway implies that when combined, these mutations will lead a loss of growth control in more than 30% of HCCs.

### M6P/IGF2 RECEPTOR, *SMAD2*, AND *SMAD4* GENES

The mannose-6-phosphate/insulin-like growth factor 2 receptor (M6P/IGF2R) is involved in the activation of transforming growth factor beta (TGF- $\beta$ ), whereas *SMAD2* and *SMAD4* genes are intracellular mediators of TGF- $\beta$ , which induces both growth inhibition and apoptotic cell death in hepatocytes.<sup>21–23</sup> After the demonstration of LOH at the M6P/IGF2R gene locus by De Souza et al.,<sup>21</sup> several reports described that the *MGP/IGF2R* gene is mutated in 18 to 33% of HCCs (Table 1). *SMAD2* and *SMAD4* genes appear to be mutated in less than 10% of these cancers.<sup>24,25</sup> In contrast, no mutation of TGF- $\beta$  receptor type II was found in HCC.<sup>24</sup> Taken together, these observations demonstrate that at least three genes involved in TGF- $\beta$ -mediated growth control are altered in HCC and that overall the TGF- $\beta$  pathway is altered in about 25% of HCCs.



**FIG. 1. Main regulatory pathways altered in human hepatocellular carcinomas.** The most frequently mutated genes of each pathway are also shown. The vertical arrowed lines connecting four pathways indicate that these pathways are related to each other and they should not be considered as independent and separate pathways of hepatocellular carcinogenesis.

## **β-CATENIN, APC, AND E-CADHERIN GENES**

The *APC* gene was initially identified in the familial adenomatous polyposis coli syndrome. Germline and somatic mutations of *APC* have been detected in colorectal cancers.<sup>26</sup> Some of these cancers display mutations in the β-catenin gene instead of the *APC* gene.<sup>27</sup> *APC* and β-catenin proteins have physical and functional interactions.<sup>28</sup> β-Catenin also forms complexes with E-cadherin.<sup>29</sup> *APC* and E-cadherin may be involved in intercellular interactions.<sup>28,29</sup> In contrast, β-catenin appears to play a role in transcriptional regulation in addition to its participation in cell-to-cell interactions.<sup>28</sup> Somatic mutations of β-catenin were observed in 19–26% of HCCs.<sup>30,31</sup> These mutations that occur at the N-terminal region of β-catenin lead to an accumulation of aberrant β-catenin proteins that stimulate the activity of a transcription factor.<sup>28,30,31</sup> Somatic *APC* mutations may be rare in HCC, but they appear to be quite frequent in hepatoblastomas.<sup>11,32–37</sup> Finally, the E-cadherin gene was shown to display frequent LOH and *de novo* methylation in HCC (Table 1). Thus, it is possible that E-cadherin function is lost in some HCCs. Taken together, these observations indicate that the “β-catenin/*APC* pathway” is altered in more than 30% of HCCs.

## **OTHER GENETIC ALTERATIONS**

As shown in Table 1, *ras* and *myc* oncogenes are not frequently mutated in human HCC. Loss of genomic imprinting and bi-allelic expression of the *IGF2* gene was shown in hepatoblastomas and in some HCCs.<sup>38–43</sup> Among other known genes, *BRCA2*, *p21*, and *p15INK4B* appear to be involved only rarely in these tumors. *MLH1* and *MSH2*, two genes involved in DNA mismatch repair, have not been studied for possible mutations in HCC (Table 1).

## **CONCLUDING REMARKS**

Recent studies clearly indicate that many genes undergo somatic aberrations (point mutations, amplifications, loss of imprinting, *de novo* methylation, etc.) in HCC. The number of aberrant genes is high, but the frequency of individual gene mutations is low. However, these mutations are not random. They tend to cluster at genes involved in important growth regulatory pathways. Even though the picture is still imperfect, our present knowledge of the molecular genetics of HCC leads us to four main pathways that are altered in HCC: the *p53* pathway involved in DNA damage response, the

*RB1* pathway involved in cell cycle control, the TGF-β pathway involved in growth inhibition and apoptosis, and the β-catenin/*APC* pathway involved in morphogenesis and signal transduction. As illustrated in Figure 1, these pathways should not be considered as independent pathways. They are most probably related to each other and may even represent individually a distinct step of hepatocellular carcinogenesis. Unfortunately, our knowledge of the order of events for the initiation and stepwise progression of HCC is still incomplete.

*Acknowledgment:* Supported by grants from TÜBİTAK, TÜBA (Turkey), and TWAS.

## **ABBREVIATIONS**

HCC	hepatocellular carcinoma
HBV	hepatitis B virus
HCV	hepatitis C virus
M6P/IGF2R	mannose-6-phosphate/insulin-like growth factor II receptor
<i>APC</i>	adenomatosis polyposis coli gene
<i>p16INK4A</i>	gene coding for a 16-kDa inhibitor of cyclin-dependent kinase 4 enzyme
<i>p15INK4B</i>	gene coding for a 15-kDa inhibitor of cyclin-dependent kinase 4 enzyme
<i>BRCA2</i>	breast cancer susceptibility gene 2
HBx	X protein of hepatitis B virus
LOH	loss of heterozygosity
<i>RB1</i>	retinoblastoma gene
pRb	protein encoded by the retinoblastoma gene
IGF2	insulin-like growth factor II
p19ARF	protein encoded by an alternatively spliced form of transcript from <i>p16INK4A</i> gene
MDM2	human homolog of mouse double mutant gene 2
p21	21-kDa cyclin-dependent kinase inhibitor protein also called CIP1
CDK4	cyclin-dependent kinase 4
E2F	a group of transcription factors regulated by the retinoblastoma family of pocket proteins
TGF-β	transforming growth factor β
<i>MLH1</i>	gene encoding a protein involved in DNA mismatch repair
<i>MSH2</i>	gene encoding for another protein involved in DNA mismatch repair

## **REFERENCES**

1. Öztürk M. Biology of hepatocellular carcinoma. In: AK Rustgi, ed. *Gastrointestinal Cancers, Biology, Diagnosis, and Therapy*. Philadelphia, PA: Lippincott-Raven, 1995, pp 511–525

2. Buendia MA. Hepatitis B viruses and hepatocellular carcinoma. *Adv Cancer Res* 1992;59:167–226
3. Idilman R, De Maria N, Colantoni A, Van Thiel DH. Pathogenesis of hepatitis B and C-induced hepatocellular carcinoma. *J Viral Hepat* 1998;5:285–299
4. Puisieux A, Ozturk M. TP53 and hepatocellular carcinoma. *Pathol Biol (Paris)* 1997;45:864–870
5. Bressac B, Kew M, Wands J, Ozturk M. Selective G to T mutations of *p53* gene in hepatocellular carcinoma from southern Africa. *Nature* 1991;350:429–431
6. Hsu IC, Metcalf RA, Sun T, et al. Mutational hotspot in the *p53* gene in human hepatocellular carcinomas. *Nature* 1991;350:427–428
7. Coursaget P, Depril N, Chabaud M, et al. High prevalence of mutations at codon 249 of the *p53* gene in hepatocellular carcinomas from Senegal. *Br J Cancer* 1993;67:1395–1397
8. Ozturk M, Puisieux A, Kew M, et al. *p53* mutation in hepatocellular carcinoma after aflatoxin exposure. *Lancet* 1991;338:1356–1359
9. Unsal H, Yakicier C, Marçais C, et al. Genetic heterogeneity of hepatocellular carcinoma. *Proc Natl Acad Sci USA* 1994;91:822–826
10. Li D, Cao Y, He L, Wang NJ, Gu JR. Aberrations of *p53* gene in human hepatocellular carcinoma from China. *Carcinogenesis* 1993;14:169–173
11. Fujimoto Y, Hampton LL, Wirth PJ, et al. Alterations of tumor suppressor genes and allelic losses in human hepatocellular carcinomas in China. *Cancer Res* 1994;54:281–285
12. Greenblatt MS, Feitelson MA, Zhu M, et al. Integrity of *p53* in hepatitis B x antigen-positive and -negative hepatocellular carcinomas. *Cancer Res* 1997;57:426–432
13. Wang XW, Forrester K, Yeh H, et al. Hepatitis B virus X protein inhibits *p53* sequence-specific DNA binding, transcriptional activity, and association with transcription factor ERCC3. *Proc Natl Acad Sci USA* 1994;91:2230–2234
14. Ueda H, Ullrich SJ, Gangemi JD, et al. Functional inactivation but not structural mutation of *p53* causes liver cancer. *Nat Genet* 1995;9:41–47
15. Elmore LW, Hancock AR, Chang SF, et al. Hepatitis B virus X protein and *p53* tumor suppressor interactions in the modulation of apoptosis. *Proc Natl Acad Sci USA* 1997;94:14707–14712
16. Greenblatt MS, Bennett WP, Hollstein M, Harris CC. Mutations in the *p53* tumor suppressor gene: clues to cancer etiology and molecular pathogenesis. *Cancer Res* 1994;54:4855–4878
17. Prives C. Signalling to *p53*: breaking the MDM2-*p53* circuit. *Cell* 1998;95:5–8
18. Reed SI. Control of the G1/S transition. *Cancer Surv* 1997;29:7–23
19. Johnson DG, Schneider-Broussard R. Role of E2F in cell cycle control and cancer. *Front Biosci* 1998;3:447–448
20. Razin A. CpG methylation, chromatin structure and gene silencing—a three-way connection. *EMBO J* 1998;17:4905–4908
21. De Souza AT, Hankins GR, Washington MK, Orton TC, Jirtle RL. *M6P/IGF2R* gene is mutated in human hepatocellular carcinomas with loss of heterozygosity. *Nat Genet* 1995;11:447–449
22. Thorgeirsson SS, Teramoto T, Factor VM. Dysregulation of apoptosis in hepatocellular carcinoma. *Semin Liver Dis* 1998;18:115–122
23. Derynck R, Zhang Y, Feng XH. Smads: Transcriptional activators of TGF-beta responses. *Cell* 1998;95:737–740
24. Kawate S, Takenoshita S, Ohwada S, et al. Mutation analysis of transforming growth factor type II receptor, smad2, and smad4 in hepatocellular carcinoma. *Int J Oncol* 1999;14:127–131
25. Yakicier MC, Irmak MB, Romano A, Kew M, Ozturk M. *Smad2* and *Smad4* gene mutations in hepatocellular carcinoma. *Oncogene* (in press).
26. Kinzler KW, Vogelstein B. Lessons from hereditary colorectal cancer. *Cell* 1996;87:159–170
27. Morin PJ, Sparks AB, Korinek V, et al. Activation of beta-catenin-Tcf signaling in colon cancer by mutations in beta-catenin or APC. *Science* 1997;275:1787–1790
28. Peifer M. Beta-catenin as oncogene: the smoking gun. *Science* 1997;275:1752–1753
29. Hirohashi S. Inactivation of the E-cadherin-mediated cell adhesion system in human cancers. *Am J Pathol* 1998;153:333–339
30. de La Coste A, Romagnolo B, Billuart P, et al. Somatic mutations of the beta-catenin gene are frequent in mouse and human hepatocellular carcinomas. *Proc Natl Acad Sci USA* 1998;95:8847–8851
31. Miyoshi Y, Iwao K, Nagasawa Y, et al. Activation of the beta-catenin gene in primary hepatocellular carcinomas by somatic alterations involving exon 3. *Cancer Res* 1998;58:2524–2547
32. Horii A, Nakatsuru S, Miyoshi Y, et al. Frequent somatic mutations of the APC gene in human pancreatic cancer. *Cancer Res* 1992;52:6696–6698
33. Kurahashi H, Takami K, Oue T, et al. Biallelic inactivation of the APC gene in hepatoblastoma. *Cancer Res* 1995;55:5007–5011
34. Giardiello FM, Petersen GM, Brensinger JD, et al. Hepatoblastoma and APC gene mutation in familial adenomatous polyposis. *Gut* 1996;39:867–869
35. Gruner BA, DeNapoli TS, Andrews W, Tomlinson G, Bowman L, Weitman SD. Hepatocellular carcinoma in children associated with Gardner syndrome or familial adenomatous polyposis. *J Pediatr Hematol Oncol* 1998;20:274–278
36. Piao Z, Choi Y, Park C, Lee WJ, Park JH, Kim H. Deletion of the *M6P/IGF2r* gene in primary hepatocellular carcinoma. *Cancer Lett* 1997;120:39–43
37. Oda H, Imai Y, Nakatsuru Y, Hata J, Ishikawa T. Somatic mutations of the *APC* gene in sporadic hepatoblastomas. *Cancer Res* 1996;56:3320–3323
38. Rainier S, Dobry CJ, Feinberg AP. Loss of imprinting in hepatoblastoma. *Cancer Res* 1995;55:1836–1838
39. Li X, Adam G, Cui H, Sandstedt B, Ohlsson R, Ekstrom TJ. Expression, promoter usage and parental imprinting status of insulin-like growth factor II (*IGF2*) in human hepatoblastoma: uncoupling of *IGF2* and *H19* imprinting. *Oncogene* 1995;11:221–229
40. Takeda S, Kondo M, Kumada T, et al. Allelic-expression imbalance of the insulin-like growth factor 2 gene in hepatocellular carcinoma and underlying disease. *Oncogene* 1996;12:1589–1592
41. Kim KS, Lee YI. Biallelic expression of the *H19* and *IGF2* genes in hepatocellular carcinoma. *Cancer Lett* 1997;119:143–148
42. Li X, Kogner P, Sandstedt B, Haas OA, Ekstrom TJ. Promoter-specific methylation and expression alterations of *igf2* and *h19* are involved in human hepatoblastoma. *Int J Cancer* 1998;75:176–180
43. Li X, Nong Z, Ekstrom C, et al. Disrupted *IGF2* promoter control by silencing of promoter P1 in human hepatocellular carcinoma. *Cancer Res* 1997;57:2048–2054
44. Yamada T, De Souza AT, Finkelstein S, Jirtle RL. Loss of the gene encoding mannose 6-phosphate/insulin-like growth factor II receptor is an early event in liver carcinogenesis. *Proc Natl Acad Sci USA* 1997;94:10351–10355
45. Murakami Y, Hayashi K, Hirohashi S, Sekiya T. Aberrations of the tumor suppressor *p53* and retinoblastoma genes in human hepatocellular carcinomas. *Cancer Res* 1991;51:5520–5525
46. Nakamura T, Iwamura Y, Kaneko M, et al. Deletions and rearrangements of the retinoblastoma gene in hepatocellular carcinoma, insulinoma and some neurogenic tumors as found in a study of 121 tumors. *Jpn J Clin Oncol* 1991;21:325–329
47. Nishida N, Fukuda Y, Kokuryu H, et al. Accumulation of allelic loss on arms of chromosomes 13q, 16q and 17p in the advanced stages of human hepatocellular carcinoma. *Int J Cancer* 1992;51:862–868

48. Zhang X, Xu HJ, Murakami Y, et al. Deletions of chromosome 13q, mutations in retinoblastoma 1, and retinoblastoma protein state in human hepatocellular carcinoma. *Cancer Res* 1994;54:4177–4182
49. Ashida K, Kishimoto Y, Nakamoto K, et al. Loss of heterozygosity of the retinoblastoma gene in liver cirrhosis accompanying hepatocellular carcinoma. *J Cancer Res Clin Oncol* 1997;123: 489–495
50. Iolascon A, Giordani L, Moretti A, Basso G, Borriello A, Della Razione F. Analysis of CDKN2A, CDKN2B, CDKN2C, and cyclin Ds gene status in hepatoblastoma. *Hepatology* 1998; 27:989–995
51. Lin YW, Chen CH, Huang GT, et al. Infrequent mutations and no methylation of CDKN2A (P16/MTS1) and CDKN2B (p15/MTS2) in hepatocellular carcinoma in Taiwan. *Eur J Cancer* 1998;34:1789–1795
52. Kita R, Nishida N, Fukuda Y, et al. Infrequent alterations of the *p16INK4A* gene in liver cancer. *Int J Cancer* 1996;67:176–180
53. Chaubert P, Gayer R, Zimmermann A, et al. Germ-line mutations of the *p16INK4(MTS1)* gene occur in a subset of patients with hepatocellular carcinoma. *Hepatology* 1997; 25: 1376–1381
54. Kim JR, Kim SY, Kim MJ, Kim JH. Alterations of *CDKN2 (MTS1/p16INK4A)* gene in paraffin-embedded tumor tissues of human stomach, lung, cervix and liver cancers. *Exp Mol Med* 1998;30:109–114
55. Furutani M, Arai S, Tanaka H, et al. Decreased expression and rare somatic mutation of the *CIP1/WAF1* gene in human hepatocellular carcinoma. *Cancer Lett* 1997;111:191–197
56. Zhang YJ, Jiang W, Chen CJ, et al. Amplification and overexpression of cyclin D1 in human hepatocellular carcinoma. *Biochem Biophys Res Commun* 1993;196:1010–1016
57. Nishida N, Fukuda Y, Komeda T, et al. Amplification and overexpression of the cyclin D1 gene in aggressive human hepatocellular carcinoma. *Cancer Res* 1994;54:3107–3110
58. Wang J, Chenivessé X, Henglein B, Brechot C. Hepatitis B virus integration in a cyclin A gene in a hepatocellular carcinoma. *Nature* 1990;343:555–557
59. Chao Y, Shih YL, Chiu JH, et al. Overexpression of cyclin A but not Skp 2 correlates with the tumor relapse of human hepatocellular carcinoma. *Cancer Res* 1998;58:985–990
60. Slagle BL, Zhou YZ, Birchmeier W, Scorsone KA. Deletion of the E-cadherin gene in hepatitis B virus-positive Chinese hepatocellular carcinomas. *Hepatology* 1993;18:757–762
61. Kanai Y, Ushijima S, Hui AM, et al. The E-cadherin gene is silenced by CpG methylation in human hepatocellular carcinomas. *Int J Cancer* 1997;71:355–359
62. Katagiri T, Nakamura Y, Miki Y. Mutations in the *BRCA2* gene in hepatocellular carcinomas. *Cancer Res* 1996;56:4575–4577
63. MacDonald GA, Greenson JK, Saito K, Cherian SP, Appelman HD, Boland CR. Microsatellite instability and loss of heterozygosity at DNA mismatch repair gene loci occurs during hepatic carcinogenesis. *Hepatology* 1998;28:90–97
64. De Benedetti VM, Welsh JA, Trivers GE, et al. *p53* is not mutated in hepatocellular carcinomas from Alaska Natives. *Cancer Epidemiol Biomarkers Prev* 1995;4:79–82
65. Shieh YS, Nguyen C, Vocal MV, Chu HW. Tumor-suppressor *p53* gene in hepatitis C and B virus-associated human hepatocellular carcinoma. *Int J Cancer* 1993;54:558–562
66. Goldblum JR, Bartos RE, Carr KA, Frank TS. Hepatitis B and alterations of the *p53* tumor suppressor gene in hepatocellular carcinoma. *Am J Surg Pathol* 1993;17:1244–1251
67. Kazachkov Y, Khaoustov V, Yoffe B, Solomon H, Klintmalm GB, Tabor E. *p53* abnormalities in hepatocellular carcinoma from United States patients: Analysis of all 11 exons. *Carcinogenesis* 1996;17:2207–2212
68. De Benedetti VM, Welsh JA, Yu MC, Bennett WP. *p53* mutations in hepatocellular carcinoma related to oral contraceptive use. *Carcinogenesis* 1996;17:145–159
69. Honda K, Sbisà E, Tullo A, et al. *p53* mutation is a poor prognostic indicator for survival in patients with hepatocellular carcinoma undergoing surgical tumour ablation. *Br J Cancer* 1998; 77:776–782
70. Soini Y, Chia SC, Bennett WP, et al. An aflatoxin-associated mutational hotspot at codon 249 in the *p53* tumor suppressor gene occurs in hepatocellular carcinomas from Mexico. *Carcinogenesis* 1996;17:1007–1012
71. Kennedy SM, Macgeogh C, Jaffe R, Spurr NK. Overexpression of the oncoprotein *p53* in primary hepatic tumors of childhood does not correlate with gene mutations. *Hum Pathol* 1994;25: 438–442
72. Kubicka S, Trautwein C, Schrem H, Tillmann H, Manns M. Low incidence of *p53* mutations in European hepatocellular carcinomas with heterogeneous mutation as a rare event. *J Hepatol* 1995;23:412–419
73. Kress S, Jahn UR, Buchmann A, Bannasch P, Schwarz M. *p53* mutations in human hepatocellular carcinomas from Germany. *Cancer Res* 1992;52:3220–3223
74. Pontisso P, Belluco C, Bertorelle R, et al. Hepatitis C virus infection associated with human hepatocellular carcinoma: Lack of correlation with *p53* abnormalities in Caucasian patients. *Cancer* 1998;83:1489–1494
75. Challen C, Lunec J, Warren W, Collier J, Bassendine MF. Analysis of the *p53* tumor suppressor gene in hepatocellular carcinomas from Britain. *Hepatology* 1992;16:1362–1366
76. Yumoto Y, Hanafusa T, Hada H, et al. Loss of heterozygosity and analysis of mutation of *p53* in hepatocellular carcinoma. *J Gastroenterol Hepatol* 1995;10:179–185
77. Debuire B, Paterlini P, Pontisso P, Basso G, May E. Analysis of the *p53* gene in European hepatocellular carcinomas and hepatoblastomas. *Oncogene* 1993;8:2303–2306
78. Scorsone KA, Zhou YZ, Butel JS, Slagle BL. *p53* mutations cluster at codon 249 in hepatitis B virus-positive hepatocellular carcinomas from China. *Cancer Res* 1992;52:1635–1638
79. Yang M, Zhou H, Kong RY, et al. Mutations at codon 249 of *p53* gene in human hepatocellular carcinomas from Tongan, China. *Mutat Res* 1997;381:25–29
80. Ng IO, Srivastava G, Chung LP, Tsang SW, Ng MM. Overexpression and point mutations of *p53* tumor suppressor gene in hepatocellular carcinomas in Hong Kong Chinese people. *Cancer* 1994;74:30–37
81. Lunn RM, Zhang YJ, Wang LY, et al. *p53* mutations, chronic hepatitis B virus infection, and aflatoxin exposure in hepatocellular carcinoma in Taiwan. *Cancer Res* 1997;57:3471–3477
82. Diamantis ID, McGandy C, Chen TJ, Liaw YF, Gudar F, Bianchi L. A new mutational hot-spot in the *p53* gene in human hepatocellular carcinoma. *J Hepatol* 1994;20:553–556
83. Sheu JC, Huang GT, Lee PH, et al. Mutation of *p53* gene in hepatocellular carcinoma in Taiwan. *Cancer Res* 1992;52:6098–6100
84. Shi CY, Phang TW, Lin Y, et al. Codon 249 mutation of the *p53* gene is a rare event in hepatocellular carcinomas from ethnic Chinese in Singapore. *Br J Cancer* 1995;72:146–149
85. Hollstein MC, Wild CP, Bleicher F, et al. *p53* mutations and aflatoxin B1 exposure in hepatocellular carcinoma patients from Thailand. *Int J Cancer* 1993;53:51–55
86. Hayashi H, Sugio K, Matsumata T, et al. Tanaka S, Sugimachi K. The mutation of codon 249 in the *p53* gene is not specific in Japanese hepatocellular carcinoma. *Liver* 1993;13:279–281
87. Nose H, Imazeki F, Ohto M, Omata M. *p53* gene mutations and 17p allelic deletions in hepatocellular carcinoma from Japan. *Cancer* 1993;72:355–360
88. Tanaka S, Toh Y, Adachi E, Matsumata T, Mori R, Sugimachi K. Tumor progression in hepatocellular carcinoma may be mediated by *p53* mutation. *Cancer Res* 1993;53:2884–2887
89. Vesey DA, Hayward NK, Cooksley WG. *p53* gene in hepatocellular carcinomas from Australia. *Cancer Detect Prev* 1994;18:123–130

90. Tsuda H, Hirohashi S, Shimosato Y, Ino Y, Yoshida T, Terada M. Low incidence of point mutation of c-Ki-ras and N-ras oncogenes in human hepatocellular carcinoma. *Jpn J Cancer Res* 1989;80:196–199
91. Tada M, Omata M, Ohto M. Analysis of ras gene mutations in human hepatic malignant tumors by polymerase chain reaction and direct sequencing. *Cancer Res* 1990;50:1121–1124
92. Stork P, Loda M, Bosari S, Wiley B, Poppenhusen K, Wolfe H. Detection of K-ras mutations in pancreatic and hepatic neoplasms by non-isotopic mismatched polymerase chain reaction. *Oncogene* 1991;6:857–862
93. Challen C, Guo K, Collier JD, Cavanagh D, Bassendine MF. Infrequent point mutations in codons 12 and 61 of ras oncogenes in human hepatocellular carcinomas. *J Hepatol* 1992;14:342–346
94. Lin SY, Chen PH, Wang CK, et al. Mutation analysis of K-ras oncogenes in gastroenterologic cancers by the amplified created restriction sites method. *Am J Clin Pathol* 1993;100:686–689
95. Leon M, Kew MC. Analysis of ras gene mutations in hepatocellular carcinoma in southern African blacks. *Anticancer Res* 1995;15:859–861
96. Zhang XK, Huang DP, Qiu DK, Chiu JF. The expression of c-myc and c-N-ras in human cirrhotic livers, hepatocellular carcinomas and liver tissue surrounding the tumors. *Oncogene* 1990;5:909–914
97. Ogata N, Kamimura T, Asakura H. Point mutation, allelic loss and increased methylation of c-Ha-ras gene in human hepatocellular carcinoma. *Hepatology* 1991;13:31–37
98. Bjersing L, Andersson C, Lithner F. Hepatocellular carcinoma in patients from northern Sweden with acute intermittent porphyria: morphology and mutations. *Cancer Epidemiol Biomarkers Prev* 1996;5:393–397
99. Tsuda H, Shimosato Y, Upton MP, et al. Retrospective study on amplification of N-myc and c-myc genes in pediatric solid tumors and its association with prognosis and tumor differentiation. *Lab Invest* 1988;59:321–327
100. Abou-Elella A, Gramlich T, Fritsch C, Gansler T. c-myc amplification in hepatocellular carcinoma predicts unfavorable prognosis. *Mod Pathol* 1996;9:95–98
101. Mares J, Polanska V, Gorgens H, et al. Oncogene amplification and expression in pediatric solid tumors. *Neoplasma* 1998; 45:123–127
102. Nishida N, Fukuda Y, Kokuryu H, et al. Role and mutational heterogeneity of the p53 gene in hepatocellular carcinoma. *Cancer Res* 1993;53:368–372
103. Oda T, Tsuda H, Scarpa A, Sakamoto M, Hirohashi S. p53 gene mutation spectrum in hepatocellular carcinoma. *Cancer Res* 1992;52:6358–6364
104. Kang YK, Kim CJ, Kim WH, Kim HO, Kang GH, Kim YI. p53 mutation and overexpression in hepatocellular carcinoma and dysplastic nodules in the liver. *Virchows Arch* 1998;432:27–32